



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2015

Oral infections: clinical and biological perspectives

Belibasakis, Georgios N ; Mylonakis, Eleftherios

Abstract: The oral cavity is a specialized ecological niche of the human body and forms a continuum with the digestive and respiratory system. The vastly diverse endogenous microbiota of the oral cavity is collectively referred to as the "oral microbiome". It has been, and continues to be, explored by numerous cultivation or culture-independent methods, yielding more than 600 individual taxa (1) . They show tropism for different microhabitats, but may also show differential associations with oral health or disease. Commensal oral microbiota live in a symbiotic relationship with the host, which is crucial for the maintenance of oral health (2) . Yet, disruption of this tight relationship by various factors will result in dysbiosis, allowing for the survival and establishment of a more virulent pathobiotic polymicrobial community that may impair the efficient immune response (3, 4) . Clinically, these events could manifest as oral infectious diseases.

DOI: <https://doi.org/10.1080/21505594.2015.1025191>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-110305>

Journal Article

Accepted Version

Originally published at:

Belibasakis, Georgios N; Mylonakis, Eleftherios (2015). Oral infections: clinical and biological perspectives. *Virulence*, 6(3):173-176.

DOI: <https://doi.org/10.1080/21505594.2015.1025191>

Oral infections: clinical and biological perspectives

Georgios N. Belibasakis^{a*}, Eleftherios Mylonakis^b

^a Section of Oral Microbiology and Immunology; Institute of Oral Biology; Center of Dental Medicine; University of Zürich; Zürich; Switzerland

^b Division of Infectious Diseases; Rhode Island Hospital; Warren Alpert Medical School of Brown University; Providence; RI; USA

* Corresponding author:

Prof. Georgios N. Belibasakis

Oral Microbiology and Immunology

Institute of Oral Biology

Center of Dental Medicine

University of Zürich

Plattenstrasse 11

8032 Zürich, Switzerland

Tel: +41-44-634-3306

Fax: +41-44-634-3090

e-mail: george.belibasakis@zsm.uzh.ch

The oral cavity is a specialized ecological niche of the human body and forms a continuum with the digestive and respiratory system. The vastly diverse endogenous microbiota of the oral cavity is collectively referred to as the “oral microbiome”. It has been, and continues to be, explored by numerous cultivation or culture-independent methods, yielding more than 600 individual taxa¹. They show tropism for different microhabitats, but may also show differential associations with oral health or disease. Commensal oral microbiota live in a symbiotic relationship with the host, which is crucial for the maintenance of oral health². Yet, disruption of this tight relationship by various factors will result in dysbiosis, allowing for the survival and establishment of a more virulent pathobiotic polymicrobial community that may impair the efficient immune response^{3,4}. Clinically, these events could manifest as an oral infectious disease.

A common oral disease of polymicrobial etiology is dental caries⁵, whose incidence and progression rate has declined in recent years due to successfully introduced population-based prevention measures^{6,7}. Progression of dental caries into the pulp space of the tooth will eventually cause pulpitis, and subsequently infection of the root canal space, leading to tooth necrosis. Expansion of the infection beyond the root canal and around the apex of the tooth may manifest as apical periodontitis, a condition associated with resorption of the periapical bone with formation of cysts and granulomas^{8,9}. This can develop unnoticed by the patient for years, or be activated with painful symptoms. While there is consensus that the interplay between the resident microbiota and the local response of the immune system are modulating the course of disease, we still fail to understand when and why these infections are activated. Zehnder and Belibasakis discuss how recent advances in molecular microbiology have given new directions in our understanding of root canal infections,

and highlight the missing links between clinical experience and experimental observations on this topic ¹⁰.

Periodontal diseases comprise a cluster of infectious inflammatory diseases that affect 10-15% of the global population in its severe form ¹¹. Gingivitis is an initial form of periodontal disease, manifesting as inflammation restricted to the superficial gingival tissue. Once this inflammation is expanded to cause destruction of the deeper periodontal (tooth-supporting) tissues, it manifests as periodontitis, the major cause of tooth loss in adults. Periodontitis compromises quality of life and its treatment poses a significant financial burden on national health systems.

Etiologically, it is the polymicrobial oral biofilms forming on the tooth surface that initiate inflammation, which may eventually become tissue-destructive. Yet, the dysbiotic host response to the biofilms is invariably a crucial determinant of the pathogenesis of periodontitis ⁴. A well-studied bacterial species frequently associated with biofilms from periodontitis-affected sites is the Gram-negative anaerobe *Porphyromonas gingivalis*. It can effectively evade the immune responses and displays invasive capacities ¹². Hence, it is featured as a “keystone” pathogen in manipulating the host’s immune response and causing dysbiosis ^{6, 13, 14}. The several virulence factors by which *P. gingivalis* can subvert the function of leukocytes, enabling it to thrive in an inflammation-rich environment are reviewed by Zenobia and Hajishengallis ¹⁵. Tissue invasion by *P. gingivalis* may also be an important virulence property of this species. To this extent Baek *et al.* show that the invasive capacity of *P. gingivalis* isolates exhibits strong positive correlations with the clinical severity of periodontitis of the subjects who harbored the isolates ¹⁶.

Among the bacteria highly associated with periodontitis is the Gram-negative *Tannerella forsythia*, a species with less well characterized virulence factors. Megson

et al have elucidated TfFuc1, a novel fucosidase expressed on its surface ¹⁷. This potential virulence factor may be linked to specific metabolic requirements of *T. forsythia* and its catalytic interactions with host proteins, potentially favoring its invasion properties. TfFuc1 is the first fucosidase to be identified in *T. forsythia*, and its biological roles warrant further investigation.

While over the years several bacterial taxa have been identified and classified as “putative” oral pathogens, several recently identified species await further characterization. As such, *Streptococcus tigurinus*, a novel species belonging to the *Streptococcus mitis* group, is discussed by Zbinden *et al.* on its association with other closely related streptococci, and whether there is evidence of its role in common oral infections ¹⁸. This is plausible given the association of *S. tigurinus* with severe medical infections, such as infective endocarditis, spondylodiscitis and meningitis ¹⁹. While there is at present only weak evidence on the involvement of *S. tigurinus* in periodontal diseases ²⁰, its presence in the oral cavity and its association with invasive infections are of high interest with regards to potential links between oral and systemic infections. At this point it should also be noted that, beyond bacterial infections, viruses may also play a role in oral pathologies. One example is human papillomavirus (HPV) infection, which has been postulated as an etiological cause of head and neck squamous cell carcinoma. Hübbers and Akgül debate on whether HPV plays also a role in the development of squamous cell carcinoma of the oral cavity ²¹.

Beyond the microbiological aspects, the immunological host response is a crucial element for the pathogenesis of oral infections. There is increasing evidence that specific genetic polymorphisms, or their combinations, confer increased susceptibility for periodontitis, or determine its progression rate ²². In a case-control study, Cavalla *et al.* monitored functional polymorphisms that result in the

transcriptional activity of the TBX21 gene, which regulates the Th-1 biased immune response²³. It was found that one of the polymorphic alleles (T) was more prevalent in chronic periodontitis, compared to gingivitis, but no association was found with presence of specific periodontal pathogens or clinical parameters of disease severity.

While the prevalence of chronic periodontitis increases by age and affects particularly the elderly population, a form of the disease of special mention is aggressive periodontitis. This is a rapidly progressive form that occurs in young individuals, who run a high risk of losing their teeth at a young age. A limited number of longitudinal studies point to predictive factors for the onset and progression of the disease being both microbial and host genetic ones. The former is backed-up by the frequent presence of *Aggregatibacter actinomycetemcomitans* JP2 clone, a Gram-negative facultative anaerobe, whereas the latter on the familial or ethnic aggregation of the disease. Hoglund Aberg *et al.* review the current knowledge on the association of *A. actinomycetemcomitans* with the risk for aggressive periodontitis, and discuss the role of its virulence factor leukotoxin in the pathogenesis of the disease²⁴. The role of the host response is further discussed by Nibali, who argues that the risk factors for aggressive periodontitis studied to-date apply only to a fragment of the cases. He reviews the evidence on the concept that genetically-driven dysbiotic changes in the subgingival biofilm microbiota may predispose to the rapid periodontal tissue destruction seen in aggressive periodontitis²⁵.

Evidently, a number of questions regarding the etiological and pathogenic mechanisms of the periodontal diseases are not fully elucidated or under question. Mendes *et al.* review the available case-control studies on the relationship between bacterial invasion of the periodontal tissues and progression periodontitis²⁶. Due to the diversity in study protocols and methodological approaches employed, no

consistent association between specific periodontal pathogens was found across studies. Hence, it is concluded that there is insufficient evidence to support or exclude the invasion of periodontal pathogens as a crucial process in the etiopathogenesis of the disease. Lopez *et al.* take an epidemiological perspective to discuss whether there is enough evidence to support our current understanding of specific periodontal pathogens as causes of periodontitis²⁷. Once again, given the strong methodological heterogeneity among studies, there are obstacles in gauging the distribution of putative periodontal pathogens across age and ethnic populations or geographic locations, with only a minimal amount of studies supporting an association between specific pathogens and progression of periodontitis. From an epidemiological perspective, it is proposed that periodontitis is better understood as an inflammatory disease resulting from the interaction of many causal components, where microbes may as well be adopted.

A special mention should be made to peri-implant infections. These have emerged following the routine use of osseointegrated dental implants in restorative dentistry. Implants provide the oral cavity with new artificial surfaces prone to the formation of biofilms, which can trigger the inflammatory destruction of the peri-implant tissue, clinically manifesting as peri-implantitis. Periodontitis and peri-implantitis feature qualitative similarities in the immune-pathological events that lead to their development, albeit the latter shows more pronounced inflammatory responses and rapid tissue destruction^{28,29}. Charalampakis and Belibasakis review the current knowledge on the microbiology of peri-implant infections gained from various methodological approaches, discussing the benefits and limitations³⁰. While there are considerable similarities in microbial profiles between periodontal and peri-

implant infections, there are also distinctive differences, which are expected to become more diversified with the increasing application of metagenomics platforms.

Research in the field of periodontal (and potentially peri-implant) infections involves clinical, *in vivo* or *in vitro* models, all of which have specific limitations, but may also complement each other. Clinical studies bring along a number of complex biological and ethical considerations, whereas animal models do not fully recapitulate the complex processes that govern human periodontitis. Hajishengallis *et al.* argue that, despite their limitations, animal models are indispensable tools for the study of the etiopathogenesis of periodontal diseases³¹. They are of particular value in testing the significance of biological mechanisms identified by *in vitro* studies, for consolidating cause-effect relationships deriving from clinical studies, and for testing novel therapeutic targets. *In vitro* experimental models tend to be over-simplistic, but on the other hand they allow for the dissection of finite biological mechanisms. *In vitro* oral biofilm models have attracted plenty of attention in recent years, particularly due to their usefulness in testing the efficacy of various anti-infective treatments^{32,33}, prior to their application in pre-clinical models or clinical trials. They are also useful in studying aspects of microbial ecology, particularly by evaluating the role of different bacteria, or their virulence factors in the function and structure of the overall biofilm community³⁴⁻³⁶. Thurnheer and Belibasakis have utilized an *in vitro* biofilm model of six oral species, in order to test if bacteria that are not part of the normal oral microbiota (*Eschericia coli*, *Staphylococcus aureus*, *Enterococcus faecails*) can efficiently colonize and grow within oral biofilms³⁷. Indeed these species were able to grow efficiently and in various structural conformations within this oral biofilm model. A derivative of this biofilm can also be used in more complex tissue-biofilm interaction models, in order to study the host response mechanisms of

relevance to periodontal disease³⁸⁻⁴⁰. Bao *et al.* established such an oral infection model resembling the periodontal pocket in a perfusion bioreactor system, which recapitulates key biological events of periodontal pathogenesis and may reduce the need for experimental animal models in the future⁴¹.

The treatment of oral infections can pose important difficulties, as these are primarily biofilm-related diseases. Biofilms are considerably more resistant against common antimicrobials treatments, compared to bacteria in their planktonic state. On the other hand, the routine use of antibiotics in common oral infections, such as periodontitis, is not ideal due to the increase in antibiotic resistance and other adverse effects. Allaker and Douglas discuss on the possibilities of non-conventional targeted approaches against biofilm-associated oral infections⁴². These include targeting a single species in the biofilm or its virulence factors, whilst maintaining the remaining of the oral microbiota. They may involve photodynamic therapy, enzyme inhibitors, enzymes and detergents, probiotics and prebiotics, as well as antimicrobial peptides and nanoparticles.

In summary, this Special Focus issue on “Oral Infections” aspires to highlight current knowledge on oral infectious diseases, while bringing forward advances in the fields of oral microbiology and immunology. Although over the years we have gathered important scientific information on periapical and periodontal infections, these still constitute important human health issues that may pose therapeutic challenges. At the same time, the more aggressive and less well-studied peri-implant infections are emerging as a future challenge for practitioners and researchers, due to the ever-increasing implementation of dental implants in practice. At this point, it is important express the notion that prevention strategies of oral infections will be on long term considerably more efficacious than their treatment. Hence, there is a need

for personalized dental healthcare and monitoring, in order to identify healthy individuals albeit at disease risk, reach an early diagnosis prior to clinical disease manifestation, and improve the prognosis of treatment outcomes. Last but not least, we should consider that the oral cavity is inseparably linked to the human body, and that the oral health may well affect the systemic health status, and vice versa.

References

1. Dewhirst FE, Chen T, Izard J, Paster BJ, Tanner AC, Yu WH, et al. The human oral microbiome. *Journal of bacteriology* 2010; 192:5002-17.
2. Devine DA, Marsh PD, Meade J. Modulation of host responses by oral commensal bacteria. *Journal of oral microbiology* 2015; 7:26941.
3. Hajishengallis G, Lamont RJ. Beyond the red complex and into more complexity: the polymicrobial synergy and dysbiosis (PSD) model of periodontal disease etiology. *Molecular oral microbiology* 2012; 27:409-19.
4. Lamont RJ, Hajishengallis G. Polymicrobial synergy and dysbiosis in inflammatory disease. *Trends in molecular medicine* 2014.
5. Simon-Soro A, Mira A. Solving the etiology of dental caries. *Trends in microbiology* 2015; 23:76-82.
6. Hajishengallis G, Lamont RJ. Breaking bad: manipulation of the host response by *Porphyromonas gingivalis*. *European journal of immunology* 2014; 44:328-38.
7. Baelum V, Heidmann J, Nyvad B. Dental caries paradigms in diagnosis and diagnostic research. *European journal of oral sciences* 2006; 114:263-77.
8. Nair PN. Pathogenesis of apical periodontitis and the causes of endodontic failures. *Critical reviews in oral biology and medicine : an official publication of the American Association of Oral Biologists* 2004; 15:348-81.

9. Belibasakis GN, Rechenberg DK, Zehnder M. The receptor activator of NF-kappaB ligand-osteoprotegerin system in pulpal and periapical disease. *International endodontic journal* 2013; 46:99-111.
10. Zehnder M, Belibasakis GN. On the dynamics of root canal infections - what we understand and what we don't. *Virulence* 2015;0.
11. Dye BA. Global periodontal disease epidemiology. *Periodontology* 2000 2012; 58:10-25.
12. Bostanci N, Belibasakis GN. *Porphyromonas gingivalis*: an invasive and evasive opportunistic oral pathogen. *FEMS microbiology letters* 2012; 333:1-9.
13. Darveau RP, Hajishengallis G, Curtis MA. *Porphyromonas gingivalis* as a potential community activist for disease. *Journal of dental research* 2012; 91:816-20.
14. Hajishengallis G, Darveau RP, Curtis MA. The keystone-pathogen hypothesis. *Nature reviews Microbiology* 2012; 10:717-25.
15. Zenobia C, Hajishengallis G. *Porphyromonas gingivalis* virulence factors involved in subversion of leukocytes and microbial dysbiosis. *Virulence* 2015;0.
16. Baek KJ, Ji S, Kim YC, Choi Y. Association of the invasion ability of *Porphyromonas gingivalis* with the severity of periodontitis. *Virulence* 2015;0.
17. Megson ZA, Koerdt A, Schuster H, Ludwig R, Janesch B, Frey A, et al. Characterization of an α -L-fucosidase from the periodontal pathogen *Tannerella forsythia*. *Virulence* 2015; 0.
18. Zbinden A, Bostanci N, Belibasakis GN. The novel species *Streptococcus tigurinus* and its association with oral infection. *Virulence* 2014;0.
19. Zbinden A, Mueller NJ, Tarr PE, Eich G, Schulthess B, Bahlmann AS, et al. *Streptococcus tigurinus*, a novel member of the *Streptococcus mitis* group, causes invasive infections. *Journal of clinical microbiology* 2012; 50:2969-73.

20. Zbinden A, Aras F, Zbinden R, Mouttet F, Schmidlin PR, Bloemberg GV, et al. Frequent detection of *Streptococcus tigurinus* in the human oral microbial flora by a specific 16S rRNA gene real-time TaqMan PCR. *BMC microbiology* 2014; 14:231.
21. Hubbers CU, Akgul B. Hpv and cancer of the oral cavity. *Virulence* 2015:0.
22. Zhang J, Sun X, Xiao L, Xie C, Xuan D, Luo G. Gene polymorphisms and periodontitis. *Periodontology* 2000 2011; 56:102-24.
23. Cavalla F, Biguetti C, Colavite P, Silveira E, Martins W, Letra A, et al. TBX21-1993T/C (rs4794067) polymorphism is associated with increased risk of chronic periodontitis and increased T-bet expression in periodontal lesions, but does not significantly impact the IFN- γ transcriptional level or the pattern of periodontopathic bacterial infection. *Virulence* 2015; 0.
24. Hoglund Aberg C, Kelk P, Johansson A. *Aggregatibacter actinomycetemcomitans*: virulence of its leukotoxin and association with aggressive periodontitis. *Virulence* 2014:0.
25. Nibali L. Aggressive Periodontitis: microbes and host response, who to blame? *Virulence* 2015:0.
26. Mendes L, Azevedo N, Felino A, Pinto MG. Relationship between invasion of the periodontium by periodontal pathogens and periodontal disease: a systematic review. *Virulence* 2015:0.
27. Lopez R, Hujoel P, Belibasakis GN. On putative periodontal pathogens: An epidemiological perspective. *Virulence* 2015; 0.
28. Belibasakis GN. Microbiological and immuno-pathological aspects of peri-implant diseases. *Archives of oral biology* 2014; 59:66-72.

29. Belibasakis GN, Charalampakis G, Bostanci N, Stadlinger B. Peri-implant infections of oral biofilm etiology. *Advances in experimental medicine and biology* 2015; 830:69-84.
30. Charalampakis G, Belibasakis GN. Microbiome of peri-implant infections: lessons from conventional, molecular and metagenomic analyses. *Virulence* 2015:0.
31. Hajishengallis G, Lamont RJ, Graves DT. The enduring importance of animal models in understanding periodontal disease. *Virulence* 2015:0.
32. Belibasakis GN, Thurnheer T. Validation of antibiotic efficacy on in vitro subgingival biofilms. *Journal of periodontology* 2014; 85:343-8.
33. Thurnheer T, Rohrer E, Belibasakis GN, Attin T, Schmidlin PR. Static biofilm removal around ultrasonic tips in vitro. *Clinical oral investigations* 2014; 18:1779-84.
34. Ammann TW, Belibasakis GN, Thurnheer T. Impact of early colonizers on in vitro subgingival biofilm formation. *PloS one* 2013; 8:e83090.
35. Ammann TW, Bostanci N, Belibasakis GN, Thurnheer T. Validation of a quantitative real-time PCR assay and comparison with fluorescence microscopy and selective agar plate counting for species-specific quantification of an in vitro subgingival biofilm model. *Journal of periodontal research* 2013; 48:517-26.
36. Bao K, Belibasakis GN, Thurnheer T, Aduse-Opoku J, Curtis MA, Bostanci N. Role of *Porphyromonas gingivalis* gingipains in multi-species biofilm formation. *BMC microbiology* 2014; 14:258.
37. Thurnheer T, Belibasakis GN. Integration of non-oral bacteria into in vitro oral biofilms. *Virulence* 2014:0.
38. Belibasakis GN, Thurnheer T, Bostanci N. Interleukin-8 responses of multi-layer gingival epithelia to subgingival biofilms: role of the "red complex" species. *PloS one* 2013; 8:e81581.

39. Belibasakis GN, Bao K, Bostanci N. Transcriptional profiling of human gingival fibroblasts in response to multi-species in vitro subgingival biofilms. *Molecular oral microbiology* 2014; 29:174-83.
40. Thurnheer T, Belibasakis GN, Bostanci N. Colonisation of gingival epithelia by subgingival biofilms in vitro: role of "red complex" bacteria. *Archives of oral biology* 2014; 59:977-86.
41. Bao K, Papadimitropoulos A, Akgul B, Belibasakis GN, Bostanci N. Establishment of an oral infection model resembling the periodontal pocket in a perfusion bioreactor system. *Virulence* 2015:0.
42. Allaker RP, Douglas CW. Non-conventional therapeutics for oral infections. *Virulence* 2015:0.